

BISMATROL- bismuth subsalicylate tablet, chewable
Preferred Pharmaceuticals Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Major Bismatrol Chewable Tablets

Active ingredient (in each tablet)

Bismuth subsalicylate 262 mg

(total salicylate 102 mg per tablet)

Purpose

Upset stomach reliever and anti-diarrheal

Uses

relieves:

- diarrhea
- heartburn
- indigestion
- nausea
- upset stomach associated with these symptoms

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. when using this product , if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- bloody or black stools
- an ulcer
- a bleeding problem

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.

Keep Out of Reach of Children

Keep Out of Reach of Children - In case of overdose, get medical help or contact a Poison Control Center immediately

Directions

- chew or dissolve in mouth
- adults and children 12 years and over: 2 tablets every 1/2 to 1 hour as needed
- do not take more than 8 doses (16 tablets) in 24 hours
- children under 12 years: ask a doctor
- drink plenty of fluids to help prevent dehydration caused by diarrhea.

Other information:

each tablet contains:

- sodium less than 1 mg
- salicylate 102 mg
- very low sodium
- avoid excessive heat (over 104°F or 40°C)
- **TAMPER EVIDENT: Do not use if individual compartments are torn or missing.**

calcium carbonate, D&C red 27 aluminum lake, flavor, magnesium stearate, mannitol, pregelatinized starch, saccharin sodium.

Principal Display Panel

MAJOR

NDC 68788-7140-3

Bismatrol Chewable Tablets




Soothing, coating action provides

fast relief for heartburn, indigestion, nausea, upset stomach and diarrhea

COMPARE TO THE ACTIVE INGREDIENT IN PEPTO-BISMOL®

Relabeled By: Preferred Pharmaceuticals Inc.

30 chewable tablets

<h1>Bismatrol Chewable Tablets</h1>			<p>Bismatrol Chewable Tablets Qty: Ins: Lot#: Bat#:</p>
<p>Generic for: Pepto-Bismol</p> <p>Active ingredient (in each tablet): Bismuth subsalicylate 262mg (each tablet contains 102 mg salicylate)</p>		<p>Bismatrol Chewable Tablets Qty: Ins: Lot#: Bat#: Prod# (NDC):</p>	<p>Bismatrol Chewable Tablets Qty: Ins: Lot#: Bat#: Prod# (NDC):</p>
<p>Pkg Size: Exp Date:</p> <p>Lot#:</p> <p>Batch#:</p> <p>Ins:</p> <p>Mfg: Major Pharm.; Livonia, MI</p> <p>Prod#:</p>	<p>Directions English</p> <p>Take <u> </u> tablet(s) every <u> </u> hours.</p>	<p>Instrucciones Español:</p> <p>Toma <u> </u> tableta(s) cada <u> </u> horas.</p>	<p>Bismatrol Chewable Tablets Qty: Ins: Insurance NDC: Lot#: Bat#:</p>
<p>Warning</p> <p>Soothing, coating action provides fast relief for heartburn, indigestion, nausea, upset stomach and diarrhea. Key's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor. Allergy alert: Contains salicylate. Do not take if you are allergic to salicylates (including aspirin) or taking other salicylate products. Do not use if you have bloody or black stools, an ulcer, a bleeding problem. Ask a doctor or pharmacist before use if you are taking any drug for anticoagulation (thinning of the blood), diabetes, gout, arthritis or if you have a fever, muscle in the stool. When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.</p>	<p>Directions English</p> <p>Take <u> </u> tablet(s) every <u> </u> hours.</p>	<p>Instrucciones Español:</p> <p>Toma <u> </u> tableta(s) cada <u> </u> horas.</p>	<p>Bismatrol Chewable Tablets Qty: Ins: Lot#: Bat#: Prod# (NDC):</p>

BISMATROL

bismuth subsalicylate tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68788-7140(NDC:0904-1315)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
STARCH, CORN (UNII: O8232NY3SJ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	17mm
Flavor		Imprint Code	GDC122
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-7140-3	5 in 1 CARTON	04/19/2018	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	04/19/2018	

Labeler - Preferred Pharmaceuticals Inc. (791119022)

Registrant - Preferred Pharmaceuticals Inc. (791119022)

Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals Inc.		791119022	RELABEL(68788-7140)

Revised: 3/2021

Preferred Pharmaceuticals Inc.